



**HUTT** maternity

Hutt Valley Maternity Care

# **Turning a Breech Baby in the Womb:** ECV (External Cephalic Version)

## Information for women

---

**What is breech?** Breech means that your baby is lying bottom or feet first in the womb (uterus). In early pregnancy breech is very common. As pregnancy continues, a baby usually turns by itself into the head first position. Between 37 and 42 weeks (term), most babies are lying head first, ready to be born.

**What is external cephalic version (ECV)?** Vaginal breech birth is more complicated than normal birth. Your obstetrician or midwife may advise trying to turn your baby to a head-first position. This technique is called external cephalic version (ECV). Gentle pressure is applied on your abdomen which helps the baby turn a somersault in the womb to lay head first. ECV increases the likelihood of having a vaginal birth.

**When can it be done?** ECV is usually tried after 37 weeks. Depending on your situation, ECV can be done right up until you give birth.

---

**Does ECV always work?** ECV is successful for about half of all women (50%). Relaxing the muscles of the womb with medication is likely to improve the chance of success. This medication will not affect the baby. You can help by relaxing your abdominal (tummy) muscles. If the baby does not want to turn, it is possible to have a second attempt on another day. If the baby does not turn after a second attempt, your obstetrician or midwife will discuss your options for birth.

**Is ECV safe for me and my baby?** ECV is generally safe and does not cause labour to begin. The baby's heart will be monitored before and after the ECV. Like any medical procedure, complications can sometimes occur. About one in 200 (0.5%) babies need to be delivered by emergency caesarean section immediately after an ECV because of bleeding from the placenta and/or changes in the baby's heartbeat.

---

ECV at Hutt Hospital is done in Delivery Suite.

**Is ECV painful?** ECV can be uncomfortable. Tell your obstetrician or midwife if you are experiencing pain so they can move their hands or stop.

ECV should not be carried out if:

- You need a caesarean section for other reasons
- You have had vaginal bleeding during the previous seven days
- The baby's heart rate tracing (also known as a ctg) is abnormal
- Your womb is not the normal pear-shape (some women have a womb which resembles a heart-shape, known as a bicornuate uterus)
- Your waters have broken before you go into labour
- You are expecting twins

## You have been booked in on:

---

- **Please arrive at Delivery Suite at 0900 hours.**
- **Please do not have anything to eat or drink from midnight the night before you come in.**

**At home after ECV.** You should telephone your Lead Maternity Carer if you have any bleeding, abdominal pain, contractions or reduced movements after ECV. There is no scientific evidence that lying down or sitting in a particular position can help your baby to turn. Always ask if you are unsure or want further information.

### **Sources and acknowledgements**

This information is based on the Royal College of Obstetricians and Gynaecologists (RCOG) guideline [External Cephalic Version and Reducing the Incidence of Breech Presentation 2006.](#)

